

**§ 520.2473**

**21 CFR Ch. I (4–1–10 Edition)**

**§ 520.2473 Tioxidazole oral dosage forms.**

**§ 520.2473a Tioxidazole granules.**

(a) *Specifications.* Each gram of granules contains 200 milligrams of tioxidazole.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Horses*—(i) *Amount.* 5 milligrams per pound of body weight as a single dose.

(ii) *Indications for use.* Removal of mature large strongyles (*Strongylus edentatus*, *S. equinus*, and *S. vulgaris*), mature ascarids (*Parascaris equorum*), mature and immature (4th larval stage) pinworms (*Oxyuris equi*), and mature small strongyles (*Triodontophorus* spp.).

(iii) *Limitations.* For administration with feed: Sprinkle required amount of granules on a small amount of the usual grain ration and mix. Prepare for each horse individually. Withholding of feed or water not necessary. Not for use in horses intended for food. The reproductive safety of tioxidazole in breeding animals has not been determined. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. It is recommended that this drug be administered with caution to sick or debilitated horses.

(2) [Reserved]

[50 FR 52772, Dec. 26, 1985; 51 FR 2693, Jan. 21, 1986, as amended at 52 FR 7832, Mar. 13, 1987]

**§ 520.2473b Tioxidazole paste.**

(a) *Specifications.* Each plastic syringe contains 6.25 grams of tioxidazole.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Horses*—(i) *Amount.* 5 milligrams of tioxidazole per pound of body weight as a single dose.

(ii) *Indications for use.* Removal of mature large strongyles (*Strongylus edentatus*, *S. equinus*, and *S. vulgaris*), mature ascarids (*Parascaris equorum*), mature and immature (4th larval stage) pinworms (*Oxyuris equi*), and mature small strongyles (*Triodontophorus* spp.).

(iii) *Limitations.* Administer orally by inserting the nozzle of the syringe through the space between front and back teeth and deposit the required

dose on the base of the tongue. Before dosing, make sure the horse's mouth contains no feed. Not for use in horses intended for food. The reproductive safety of tioxidazole in breeding animals has not been determined. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. It is recommended that this drug be administered with caution to sick or debilitated horses.

(2) [Reserved]

[52 FR 43059, Nov. 9, 1987]

**§ 520.2475 Toceranib.**

(a) *Specifications.* Each tablet contains 10, 15, or 50 milligrams (mg) toceranib as toceranib phosphate.

(b) *Sponsor.* See No. 000009 in § 510.600 of this chapter.

(c) *Conditions of use*—(1) *Dogs*—(i) *Amount.* Administer an initial dose of 3.25 mg per kilogram (1.48 mg per pound) body weight, orally every other day.

(ii) *Indications for use.* For the treatment of Patnaik grade II or III, recurrent, cutaneous mast cell tumors with or without regional lymph node involvement.

(iii) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[74 FR 28875, June 18, 2009]

**§ 520.2483 Triamcinolone.**

(a) *Specifications.*—(1) Each tablet contains 0.5 milligram (mg) or 1.5 mg triamcinolone acetonide.

(2) Each 15 grams of powder contains 10 mg triamcinolone acetonide.

(b) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(c) *Special considerations.* See § 510.410 of this chapter.

(d) *Conditions of use*—(1) *Dogs and cats.* Use tablets described in paragraph (a)(1) of this section as follows:

(i) *Amount.* Administer 0.05 mg per pound (lb) of body weight daily by mouth; up to 0.1 mg per pound (lb) of body weight daily, if response to the smaller dose is inadequate. Therapy may be initiated with a single injection of triamcinolone acetonide suspension as in § 522.2483 of this chapter, in which case triamcinolone acetonide